## **Species Notes for**

# Pinyon Mouse (Peromyscus truei):

# California Wildlife Habitat Relationships (CWHR) System Level II Model Prototype



William F. Laudenslayer, Jr. (author)<sup>1</sup> and Monica D. Parisi (editor)<sup>2</sup>

California Department of Fish and Game California Interagency Wildlife Task Group

October, 2007

<sup>&</sup>lt;sup>1</sup> United States Forest Service, Pacific Southwest Research Station (ret.)

<sup>&</sup>lt;sup>2</sup> California Department of Fish and Game, Biogeographic Data Branch

#### **PREFACE**

This document is part of the California Wildlife Habitat Relationships (CWHR) System, operated and maintained by the California Department of Fish and Game (CDFG) in cooperation with the California Interagency Wildlife Task Group (CIWTG). The information will be useful for environmental assessments and wildlife habitat management. For more information on the CWHR System and all of its components, please see <a href="http://www.dfg.ca.gov/biogeodata/cwhr/">http://www.dfg.ca.gov/biogeodata/cwhr/</a>.

Notes such as these were prepared for 32 species by the US Forest Service Pacific Southwest Research Station as part of a 2000/2001 contract with CDFG. Each is part of a prototypical "Level II" model for a species. As compared with the "Level I" or matrix models initially available in the CWHR System, "Level II" models incorporate spatial issues such as size of a habitat patch and distance between suitable habitat patches.

The notes are divided into three major sections. First, "Distribution, Seasonality and Habitats" represents information in the existing Geographic Information System (GIS) range data and in the Level I matrix model for a species. There is a vector-based GIS layer of geographic range and seasonality for each species in CWHR as well as a matrix containing all suitability ratings – High (H), Medium (M), Low (L) or Unsuitable (-) – by habitat (e.g. BOW or Blue Oak Woodland), stage (e.g. 4P or small tree, open canopy) and life requisite (reproduction, cover, or feeding.). Tools such as "Bioview" within the CWHR software will return these suitability ratings for a species to a user-supplied data set containing habitats and either stages (e.g. 4P) or stage values (e.g. trees of 16.0 average diameter at breast height in a stand of 30% canopy closure).

Second, "Required Attributes of Suitable Habitat Patches" represents spatially-explicit requirements of a species. The information here builds upon what is known about habitat patch size and the most critical attributes of a habitat patch needed by an individual of the species. Applications such as "GRABS", which stands for "Grouping Resources Algorithm for Biological Data Sets", will "clump" pixels of a user-supplied raster-based GIS data set representing patches of a suitable habitat and stage for a species. It will calculate area, perimeter, and complexity within each patch and analyze its outside edge for juxtaposition with other habitats and stages of interest. Many of the attributes are what were once called "elements" in the CWHR model.

Third, "Spatial Habitat Requirements for Persistence of Population" represents estimates of the amount of habitat needed to maintain a population of a species. This may be considered the starting point for a "Level III" CWHR model, which would take into account spatial issues as well as a number of population parameters not yet incorporated into CWHR. Such information is included for most, but not all, Level II-modeled species.

### **Distribution, Seasonality and Habitats**

Model Parameter	Threshold Value(s) for Species
Biogeographic Range and Seasonality range of the species, by season, in the state	Species is found in forested and shrub habitats throughout much of California except the Central Valley, most of the Mojave Desert, high elevations in the Sierra Nevada, and the coast of southern California. In warmer climates (i.e. at lower elevations) these mice are active much of the year, but in colder climates they generally winter in dens.
habitats rated in the California Wildlife Habitat Relationships (CWHR) System as high (H), medium (M), or low (L) suitability for reproduction, cover, or feeding	Species finds suitability (H>L) for reproduction, cover and/or feeding in some or all stages of: Alkali Desert Scrub, Alpine Dwarf-Shrub, Annual Grassland, Bitterbrush, Chamise-Redshank Chaparral, Coastal Scrub, Desert Riparian, Desert Wash, Douglas Fir, Eastside Pine, Eucalyptus, Juniper, Klamath, Mixed Conifer, Low Sage, Mixed Chaparral, Montane Chaparral, Montane Hardwood, Montane Hardwood-Conifer, Montane Riparian, Palm Oasis, Perennial Grassland, Pinyon-Juniper, Ponderosa Pine, Redwood, Sagebrush, Sierran Mixed Conifer, Valley Foothill Riparian, Wet Meadow, and White Fir.
_	Water enhances suitability. This mouse is generally found in rather xeric conditions. Water is acquired through prey items and free water such as dew.

## **Required Attributes of Suitable Habitat Patches**

Model Parameter	Threshold Value(s) for Species
Patch Size	0.25 acre (L)
L = low suitability. This is	5 acres (H)
the minimum patch size for	
persistence of an individual.	
H = high suitability. Above	
this patch size, area alone	
does not increase habitat	
suitability for an individual.	

requirements for a transition between two life form types – tree/shrub, tree/grass, tree/water, tree/agricultural, shrub/grass, shrub/water,	Edges are not required by this species.
shrub/agricultural, grass/water, grass/agricultural, or water/agricultural	
Structural Habitat Attributes requirements for live vegetation, dead or decadent vegetation, vegetation residues, physical features, or human-made features	A shrub layer or herb layer is required for reproduction, cover and feeding. A relatively continuous cover of herbs and shrubs within a habitat patch is desirable. Lithic, rock, snags, and all sizes of slash and logs enhance the value of habitat.
Food vegetative or animal diet requirements	Preferred diet items include graminoids, herbs, seeds, acorns, berries, fungi and invertebrates.

### **Spatial Habitat Requirements for Persistence of Population**

Lowest suitability = 200 acres, if suitable patches are of a minimum size (see above) and are a maximum of 10 meters apart

Highest suitability = greater than 200 acres, if suitable patches are of a minimum size (see above), and are less than 5 meters apart